

year. The plan has infinite possibilities in the safeguarding of a citrus investment, and no doubt in the future mortgagees will insist that a crop be insured and when the time comes that the companies will consider tree insurance citrus property will take its place as security along with other real estate securities; stabilizing the industry; possibly making Government farm-loan funds more available.—*Florida Grower*, Nov. 28, 1921.

ARCADIA, Nov. 21.—The first frost insurance policy ever written in this city was written here Monday in favor of the Arcadia Citrus Exchange branch of the county organization for the amount of \$75,000 to cover this year's crop from frost or freeze. The premium is 10 per cent of the gross amount of the insurance carried and will cost the grower members of the exchange \$7,500, which is divided pro rata on the box basis, and will cost the 250 members 3 cents per box, basing the estimate on a normal crop.—*Tampa (Fla.) Tribune*, Nov. 28, 1921.

ATMOSPHERIC ELECTRICITY AND THE MOVEMENT OF DEPRESSIONS.¹

By J. LACOSTE.

[Abstracted from *Comptes Rendus*, Nov. 7, 1921, pp. 843-845.]

Studies made during the summer of 1921 at the Geophysical Institute of Strasbourg, on radiogoniometry, have led to interesting meteorological relations between the direction of maximum intensity of "strays" and the movement and shape of depressions. The author has stated four tentative laws concerning this relation:

1. In the case of a well-defined circular depression, the maximum intensity of strays is observed in the direction of the southern or southeastern part of the depression: the change of this direction of maximum intensity serves to follow the movement of the depression.

2. In the case of an elliptical depression, less well marked, the maximum intensity, as in the first case, is observed to come from the southern or southeastern parts. This case is less satisfactory than the first.

3. In the case of secondary depressions, or the barometric fluctuations along squall fronts, the maximum is difficult to determine.

4. A squall close at hand gives violent strays.

These observations have permitted the author to follow the movements of known depressions, and also to forecast the arrival of those as yet unseen on the weather map. He also found a striking relation between observations of maxima in 1920 and the corresponding barometric distribution. This, he believes, points the way to a new viewpoint in the forecasting of weather.—*C. L. M.*

551.509.6 (048)

ARTIFICIAL PRODUCTION OF RAIN.

By DR. HAROLD JEFFREYS.

[Reprinted from *Nature*, London, Nov. 3, 1921, pp. 312-314.]

In an article in the *Times* of October 17 an account is given of the achievements of Mr. Charles M. Hatfield in producing rain. The method used is not described in any detail. A tank filled with certain unspecified "chemicals" was exposed at a height of 25 feet above the ground, and it is claimed that this had the effect of producing 8 inches of rain in three months at Medicine Hat, 22 miles away. The theory of the method is that the apparatus draws clouds from other parts to the Medicine Hat district and causes them to precipitate their moisture there. No direct observations of the motions of clouds are mentioned in confirmation of this theory, though they should not have been difficult to obtain.

The official raingage at Medicine Hat during May, June, and July, the period of the contract, recorded 4.8 inches, which was 1.3 inches below the normal for the station for those months. Further comment on the success of the experiments is unnecessary.

The financial side of Mr. Hatfield's contract with the United States Agricultural Association of Medicine Hat is interesting, for the association was apparently prepared to pay Mr. Hatfield as if 8 inches of rain had fallen. Still more interesting is the fact that he was promised \$4,000 for 4 inches and \$6,000 for 6 inches. Since the normal rainfall is 6.1 inches Mr. Hatfield would have been much more likely than not to make a substantial profit even if he had done nothing at all.

It may be mentioned that at Calgary, Alberta, the rainfall was 3 inches below normal; at Edmonton, it was 3.1 inches above; and at Qu'Appelle, Saskatchewan, 300 miles to the east, it was 3.85 inches above normal.

It is also stated that at Los Angeles, in the first four months of 1905, Mr. Hatfield guaranteed 18 inches of rain, and that his own raingage showed 29.49 inches. If this is correct the rainfall must have been extremely local, for the official raingage at Los Angeles in those months, showed only 14.98 inches. Still, this was 4.4 inches above normal. At San Diego, however, which is 200 miles away, the excess was 4.6 inches, and it appears likely that the abnormality at both stations was due to more widespread causes than Mr. Hatfield's chemicals.

Attempts have on many previous occasions been made to produce rain by artificial means, but the results have been uniformly unsuccessful. The reason is not difficult to see. To make the water vapor in the air condense it is necessary to cool the air in some way to a temperature below the dew point. This may be done in two ways. One may cool the air directly, for instance by the evaporation of liquid carbon dioxide or liquid air. This certainly would produce a little condensation; the fatal objection to it is that it would be thousands of times cheaper to distil sea water. The other method is to raise the air. The pressure decreases with height, and to reduce the pressure on a particular mass of air is known to cool it. The difficulty is to raise it enough. To produce an inch of rain over an area of 100 square miles requires the condensation of 6,000,000 tons of vapor, and to achieve this some hundreds of millions of tons of air must be lifted up. The distance it must be raised depends on how nearly saturated it was originally, but it could not be less than a kilometer in ordinary fine weather conditions. We have no source of energy at our command great enough to achieve this.

It is often suggested that rain may be produced by exploding shells or otherwise agitating the air. The action is compared with that of a trigger, a large amount of energy being released by a small effort. An essential feature is, however, overlooked. For a trigger to work, there must be a large supply of potential energy only awaiting release. Precipitation from partially saturated air would require an actual supply of new energy. Therefore a trigger action can not produce precipitation.

Apropos of the above, the following account of rain-maker Hatfield's exploit in the State of Washington is not without interest:

The people of Washington have taken a lively interest in the efforts of a professional rainmaker near Moses Lake, who was engaged by the Commercial Club of Ephrata to arrange for enough precipitation to insure bountiful crops this summer. Hundreds of visitors, bitten by

¹ Sur la relation existant entre les directions des dépressions et les directions des maxima des parasites atmosphériques.

the curiosity, have gazed upon the curious tower from which mysterious gases were supposed to flow into the atmosphere. As a sporting proposition the attempt of the rain maker was not without public interest; if he produced rain he was to receive adequate remuneration, but if he failed he was to have nothing but the inexpensive good wishes of the community. Now the rainmaker has gone and almost immediately the skies open up and discharge their heavy load of moisture. Skep-

tical persons may refuse to see any connection between the mysterious "forces" loosed by the rainmaker and the cloudbursts in eastern Washington, but the miracle man, it appears, is able to recognize the storms as his own. Especially, he avers, does the rainfall at Ephrata have his private brand, although its arrival was somewhat late.—** * * Post Intelligencer*, Seattle, Wash., July 17, 1920. Reprinted from *Bulletin American Meteorological Soc.*, Aug. 1920, pp. 80-82.

BIBLIOGRAPHY.

RECENT ADDITIONS TO THE WEATHER BUREAU LIBRARY.

C. FITZHUGH TALMAN, Professor in Charge of Library.

The following have been selected from among the titles of books recently received as representing those most likely to be useful to Weather Bureau officials in their meteorological work and studies:

Fédération aéronautique internationale.

Table de conversion des pressions barométriques en altitudes. Paris. 1920. 8. p. 17½ cm.

France. Service hydrométrique du bassin de la Seine.

Résumé des observations centralisées. Versailles. 1919. 19 p. 31 cm.

Great Britain. Meteorological office.

British rainfall. 1920. London. 1921. fronts. illus. plates. maps. xxviii, 285 p. 21½ cm.

Great Britain. Meteorological office.

Forecast code for the abbreviation of weather forecasts transmitted by telegraphy or radiotelegraphy. London. 1921. 18 p. diagr. 24½ cm. (M. O. 244.)

Great Britain. Meteorological office.

Marine observer's handbook. 3d ed. London. 1921. iv, 99 p. illus. plates. 24½ cm. (M. O. 218.)

Great Britain. Meteorological office.

Observer's handbook. 1921 ed. London. 1921. xxx, 110, 10 p. illus. plates. 24½ cm. (M. O. 191.)

Italy. Consiglio superiore delle acque pubbliche.

Annali. Utilizzazione delle acque—Irrigazione—Concessioni di acque pubbliche. Anno 1921. v. 3. fasc. 1-2. Roma. 1921. 25½ cm.

Jefferson, Mark.

Rainfall of Chile. New York. 1921. 32 p. illus. maps. 19 cm. (American geographical society. Research series no. 7.)

Sifontes, Ernesto.

El método de Boussingault y la temperatura media. Anual 1-3. Ciudad Bolívar. 1920-1921. sheets. (Climatología y meteorología tropical.)

Smith, L. A. Brooke.

Weather forecasting in the eastern North Atlantic and home waters for seamen. London. 1921. 23 p. plates. 25 cm. (Great Britain Meteorological office. M. O. 246.)

Southport (England.) Fernley observatory.

Report and results of meteorological observations. By Joseph Baxendell. 1920. Southport. 1921. 32 p. 25 cm.

Sumner, P. H.

Design and stability of streamline kite balloons, with useful tables, aeronautical and mechanical formule. London. 1920. viii, 146 p. illus. fronts. plate. 22 cm.

RECENT PAPERS BEARING ON METEOROLOGY AND SEISMOLOGY.

C. F. TALMAN, PROFESSOR IN CHARGE OF LIBRARY.

The following titles have been selected from the contents of the periodicals and serials recently received in the Library of the Weather Bureau. The titles selected are of papers and other communications bearing on meteorology and cognate branches of science. This is not a complete index of all the journals from which it

has been compiled. It shows only the articles that appear to the compiler likely to be of particular interest in connection with the work of the Weather Bureau.

Aerial age weekly. N. Y. v. 14. Dec. 5, 1921.

Hersey, M. D. Aneroid investigations in Germany. p. 296-297.

Aérophile. Paris. 29 année. 1-15 octobre, 1921.

Roux, G. La météorologie et les grandes épreuves de distance. p. 291-295.

American academy of arts and sciences. Proceedings. Boston. v. 56. Sept., 1921.

Ward, Robert DeC. Walter Gould Davis. (1851-1919) p. 402-405. [Obituary.]

American society of heating and ventilating engineers. Journal. N. Y. v. 57. Nov., 1921.

Stacey, A. E., jr. Notes on the relation of atmospheric conditions to chemical processes. p. 777-781.

Annalen der Hydrographie und maritimen Meteorologie. Berlin. 49 Jahrg. 1921.

Bingel, Fr. Turbulenz und Vertikalströme in der freien Atmosphäre. Teil 1-2. H. 7-8. p. 209-220; 248-257.

Hanzlik, St. Luftdruck und Wärmeschaukel in der elfjährigen Klimaperiode. H. 8. p. 263-267.

Spitaler, Rudolf. Eine neue Theorie der täglichen Luftdruckschwankung sowie der Ebbe und Flut überhaupt. H. 7. p. 220-231.

Annalen der Physik. Leipzig. Bd. 65. 1921.

Hochschwender, E. Über das Zerblasen von Wassertropfen im Luftstrom und die Wasserfalltheorie der Gewitter. p. 629-639.

Annali del Consiglio superiore delle acque. Roma. v. 3. fasc. 2. 1921.

Eredia, Filippo. Le abbondanti piogge del settembre 1920 sul Veneto orientale. p. 51-72.

Montanari, Vittorio. Nubifragi e laghi artificiali. p. 73-77.

Astronomie. Paris. 35 année. Nov., 1921.

Flammarion, Camille. Un été exceptionnel. Le poids de la chaleur. p. 409-415.

Ecology. Brooklyn. v. 2. July, 1921.

Clements, Frederic E. Drouth periods and climatic cycles. p. 181-188.

Engineering news-record. N. Y. v. 87. 1921.

Drought in the British Isles. p. 820. (Nov. 17.)

Hazen, Allen. Frequency of high rates of rainfall; a probability-method study of data for the eastern and middle western states in Morgan's Miami technical reports, part V, and Meyer's "Elements of hydrology." p. 858-860. (Nov. 24.)

France. Académie des sciences. Comptes rendus. Paris. T. 173. 1921.

Mascart, Jean. Sur l'établissement des moyennes en météorologie. p. 94-96. (11 juil.)

Schereschewsky, Ph. Les fondements de la classification rationnelle des nuages. p. 96-98. (11 juil.)

Nodon, Albert. Recherches expérimentales sur des relations entre les phénomènes électriques terrestres, l'état de l'atmosphère et les foyers solaires. p. 722-724. (24 oct.)

Brazier, C. E. La variation de la vitesse ascensionnelle des ballons pilotes avec l'altitude. p. 756-758. (2 nov.)

Lacoste, J. Sur la relation existant entre les directions des dépressions et les directions des maximums des parasites atmosphériques. p. 843-845. (7 nov.)

Guilbert, Gabriel. Sur la formation de la pluie et l'origine des cirrus. p. 999-1001. (21 nov.)

Kerforné, F., & Milon, Y. Observations sur deux sismes récents du massif armoricain. p. 995-997. (21 nov.)

Lacoste, J. Contribution à la prévision du temps et en particulier des orages par les visées de ballons-pilotes. p. 997-999. (21 nov.)

Schereschewsky, Ph., & Wehrli, Ph. Sur le mouvement des noyaux de variations de pression. P. 1001-1004. (21 nov.)